

Stream water nutrient and organic carbon exports from tropical headwater catchments at a soil degradation gradient

John W. Recha · Johannes Lehmann ·
M. Todd Walter · Alice Pell · Louis Verchot ·
Mark Johnson

Received: 23 July 2012 / Accepted: 15 January 2013 / Published online: 31 January 2013
© Springer Science+Business Media Dordrecht 2013

Abstract Carbon and nutrient losses were quantified from four small headwater catchments in western Kenya in the year 2008. They include a forested catchment and three catchments under maize continuously cultivated for 5, 10 and 50 years following forest conversion. The C isotopic composition of dissolved organic C (DOC) in stream discharge suggested that soil organic C (SOC) derived from the original forest rather than OC from maize may have contributed to a large extent to

watershed OC losses, even 50 years after the forest was removed. Flow-weighted stream water concentrations of DOC and coarse particulate OC, all N species, total P, K and Na significantly ($P < 0.05$) increased in streams after forest conversion and long-term cultivation. Solute concentrations increased despite the fact that soil contents decreased and total water flow increased indicating mobilization of C and N, P and K from soil with progressing cultivation. In contrast, Ca and Mg concentrations in stream water did not systematically change after deforestation and cultivation, and may be controlled by geochemical weathering rather than by changing water flow paths or topsoil contents. All OC and nutrient exports increased with longer cultivation over decadal time scales ($P < 0.05$) to the same or greater extent than through deforestation and the first years of cultivation. Fluvial OC and total N losses were 2 and 21 % of total SOC and total N decline, respectively, in the top 0.1 m over 50 years. Fluvial OC losses therefore played a minor role, and SOC losses were mainly a result of microbial mineralization. Resulting total N losses by stream discharge, however, were large with $31 \text{ kg ha}^{-1} \text{ year}^{-1}$ after 50 years of continuous cropping in comparison to fertilization of $40 \text{ kg N ha}^{-1} \text{ year}^{-1}$. Most (91 %) of the N losses occurred as NO_3^- . In contrast, P losses by stream discharge were negligible in comparison to plant uptake. Water losses should be managed to reduce soil fertility declines especially through large N export from agricultural headwater catchments. However, stream concentrations of both P (0.01–0.15 mg L^{-1})

J. W. Recha · J. Lehmann (✉)
Department of Crop and Soil Sciences, Cornell
University, Ithaca, NY, USA
e-mail: CL273@cornell.edu

M. T. Walter
Department of Biological and Environmental
Engineering, Cornell University, Ithaca, NY, USA

A. Pell
Department of Animal Sciences, Cornell University,
Ithaca, NY, USA

L. Verchot
Center for International Forestry Research, Bogor,
Indonesia

M. Johnson
Institute for Resources, Environment and Sustainability,
Vancouver, Canada

M. Johnson
Department of Earth and Ocean Sciences, University
of British Columbia, Vancouver, Canada

and N ($0.4\text{--}4.8\text{ mg L}^{-1}$) were moderate or low with respect to possible consequence for human health and not responsible for eutrophication observed in Lake Victoria.

Keywords Carbon · Cultivation · Degradation gradient · Forest conversion · Headwater catchment · Nutrients

Introduction

The impact of land use change on nutrient and soil organic carbon (SOC) dynamics in the tropics is particularly severe since soil degradation is more rapid than in temperate zones (Spaans et al. 1989; Malmer and Grip 1990; Grip et al. 2004; Hartemink et al. 2008). Continuous cultivation and land tillage cause rapid loss of SOC (Mann 1986; Davidson and Ackerman 1993; Tilman et al. 2002; McLauchlan 2006; Solomon et al. 2007) due to the disruption of the physical, biochemical, and chemical mechanisms of soil organic matter (SOM) stabilization. However, the reduction in SOC stocks may not be solely attributed to increased mineralization losses to CO_2 , but could include a significant, yet poorly quantified proportion of stream water losses (Davidson and Ackerman 1993). On a basin and headwater catchment scale of forests, stream water OC losses appear to be a minor fraction of total SOC losses compared to mineralization of CO_2 (Richey et al. 2002; Johnson et al. 2006b). However, the importance of fluvial OC losses during the first years of conversion from forests to cultivation, which are the most important in terms of SOC losses (Davidson and Ackerman 1993) have not been well quantified. While mineralization losses typically decrease with greater period of cultivation (Kimetu et al. 2009), the stream water discharge usually increases (Williams et al. 1997; Germer et al. 2010; Recha et al. 2012), potentially increasing the importance of fluvial OC losses as cultivation progresses.

Soil organic matter decline leads to rapid nutrient release of N and P as well as reduced cation exchange capacity resulting in diminished nutrient retention of the soil (Lal 2006). However, surprisingly little information is available on how soil degradation following land use conversion affects nutrient losses from watersheds. On a plot level, information on accelerated decrease in soil nutrient contents resulting from various types of agricultural practices has been

gathered in both temperate (Nair and Graetz 2004; Schipper et al. 2007) and tropical regions (Pandey et al. 2010). However, studies on nutrient losses from entire catchments have mostly been confined to forests rather than to different agricultural watersheds with increasing period of cultivation. Studies from Malaysia and the Amazon examined impacts of disturbances (Zulkifli 1990), pasture (Thomas et al. 2004) and forest clearing (Lesack 1993; Malmer 1996; Williams and Melack 1997; Williams et al. 1997). It is not clear, to what extent continuous cultivation and accompanying soil degradation over time affect fluvial nutrient losses in addition to the effects of loss of forest cover, and how these fluvial losses compare to nutrient input and crop uptake.

Not only the relationship between agricultural nutrient input and fluvial export is unclear, but also what these exports mean for pollution of waterways. Lake Victoria, the largest tropical lake in the world, is significantly contaminated with DOC and nutrients, notably P, posing a threat to people, the fish industry and biodiversity (Verschuren et al. 2002). It is widely believed that the contamination originates from landuse change induced by agriculture (Scheren et al. 2000; Hecky et al. 2010). This is a common attribution and non-point source pollution with N and P by agriculture is indeed observed in many parts of the world (Carpenter et al. 1998). Yet, fertilizer use in tropical smallholder agriculture is typically much lower than in industrialized countries where detailed assessments are available. No data are available that investigate the nutrient and OC loads of agricultural headwaters in Africa and the contribution that deforestation by burning and progressive cultivation would make.

The dynamics of nutrient and C losses over the course of the year can provide insight into the mechanism and pathways of nutrient and C in watersheds (Markewitz et al. 2004; Johnson et al. 2006a). Typically, we expect a dilution of nutrient and C concentrations during the rainy season (Likens and Bormann 1995). However, under certain circumstances of highly weathered soils, concentrations may increase during the wet season as a result of greater surficial flow paths that mobilize solutes from nutrient and C-rich topsoils (Markewitz et al. 2004). It is not clear whether forest conversion and soil cultivation may change the relationship between water and nutrient concentrations and how these dynamics

are modified in tropical soils rich in weatherable minerals (Davidson et al. 2004; Yusop et al. 2006) with high exchangeable base cation contents (Kimetu et al. 2008). The fluvial DOC losses may be a significant fraction of C losses and of total solute losses in watersheds (Selva et al. 2007). In forest ecosystems, the majority of the fluvial organic C losses occurred as DOC rather than coarser organic C fractions (Johnson et al. 2006b; Selva et al. 2007). Despite the lower importance of particulate organic C, most of the organic C was mobilized from the topsoil litter layer (Johnson et al. 2006a, b). Whether the same is true for agricultural soils, or whether subsoil organic C may gain in importance for soils depleted in SOC by long-term cultivation is not clear.

The purpose of this study was to quantify the fluvial nutrient and OC losses from headwater catchments separating the effects of soil degradation and concomitant losses of soil organic matter contents from those of forest clearing in the upper catchments of the Lake Victoria basin. It is hypothesized that (1) stream water OC losses increase in importance with length of cultivation; (2) long-term cultivation is as or more important than deforestation and short-term cultivation for increasing nutrient and OC losses; and (3) headwaters with small holder agriculture are not important sources of N and P to rivers and lakes. The effects of continuous agricultural land use on nutrients were investigated by comparing three headwater catchments following forest conversion to continuous maize production for either 5, 10, or 50 years in Western Kenya with a catchment that has remained forested.

Methods

The study site

The field measurements were done in Kapchorwa within Nandi County in Kenya, which is located in Eastern Africa. The site is located 60 km northeast of Lake Victoria at longitude 35°0'0" E and latitude 0°10'0"N (Recha et al. 2012). The region belongs to the sub-humid ecological zone characterized by a bimodal rainy season with a mean annual precipitation of 2,000 mm (Awiti et al. 2008). The “long rain season” is from April to June (~1,200 mm) and the “short rain season” from August to October (~800 mm).

The site has a mean elevation of 1,800 m above sea level, a maximum daily temperature of 26 °C and a minimum of 11 °C (Glenday 2006).

The Kakamega-Nandi forest in Western Kenya is the country's only remaining tropical rain forest. Massive deforestation has taken place to create land for settlement and farming. About 16 % of forest cover was lost between 1986 and 2001 (Awiti et al. 2008). The forest forms the easternmost relic of the Guinean-Congolian rainforest belt, which once spanned from East to West Africa. The area around this forest is among the most densely populated rural areas in the world. It had a population density of 778 persons per km² in 2009 compared to 73 persons per square kilometer for the entire country (Kenya National Bureau of Statistics 2010). Consequently the forest is under high anthropogenic pressure, which is mirrored by the decreasing natural forest cover and intensive cultivation (deGraffenried and Shepherd 2009; Swallow et al. 2009). Past deforestation rates in the Kakamega-Nandi forest indicated a decrease of forest area and an increase in the fragmentation of natural, old-growth forest (Lung and Schaab 2006).

Soils in the Kapchorwa catchment are kaolinitic Acrisols (FAO-UNESCO-ISRIC 1988), which are classified as Ultisols in the US soil taxonomy (Soil Survey Staff 2003). The parent material of these soils is principally granite, with some inclusions of Precambrian gneisses, which supports Luvisols (Werner et al. 2007) and other undifferentiated basement system rocks at higher elevations (Jaetzold and Schmidt 1983). Soils in the catchment have 45–49 % clay, 15–25 % silt, and 26–40 % sand (Kimetu et al. 2008).

The forest section of the Kapchorwa catchment is part of the Kakamega-Nandi forest composed of tropical rainforest species. It is largely an indigenous forest with a 30-m closed canopy dominated by evergreen hardwood species. The most common species are *Funtumia africana*, *Ficus* spp, *Croton* spp, and *Celtis* spp (Glenday 2006). Other species include *Aningeria altissima* (A. Chev.), *Milicia excelsa* (Welw.) C.C. Berg, *Antiaris toxicaria* (Lesch.) and *Chrysophyllum albidum* (G. Don), *Olea capensis* (L.) and *Croton megalocarpus* (Hutch.) (Kinyangi 2008). The above and below ground net primary production of trees in a tropical forest is estimated at 15.2 Mg ha⁻¹ year⁻¹ (Hertel et al. 2009). The agricultural catchments have maize as the sole crop, and

have been under maize cultivation since conversion from forest cover. The maize grain yields without fertilizer input in the 5-, 10- and 50-year old agricultural catchments are 6.5, 5.5 and 2.5 Mg ha⁻¹ year⁻¹, respectively (Ngoze et al. 2008).

Hydrologic instrumentation and fieldwork

In order to identify the headwater catchments, the time when native forests were converted to agriculture was determined from historical community settlement patterns over the last century (Bleher et al. 2006; Recha et al. 2012). Specific years of conversion were verified from data available from records of the Kenyan government from the Department of Forests, the Ministry of Agriculture, as well as from interviews with officials of local institutions and from county council records. Within each site, population settlement patterns were distinct and newly acquired fields were excised from sections of the native forest for agriculture. All four headwater catchments are located within an area of 6 km² (Fig. 1) and represent a soil degradation gradient that corresponds to years under maize cultivation that has been used in several other studies (Kimetu et al. 2008, 2009; Ngoze et al. 2008; Kimetu and Lehmann 2010; Recha et al. 2012). Such chronosequences substitute time for space and have to be carefully selected to assure similar properties before the change (Huggett 1998). Also hydrological differences between catchments have to be considered (Elsenbeer, 2001; Johnson et al. 2006b) and only clear trends across the entire set of the four catchments are interpreted here.


Hydrologic instrumentation was installed in 2006 (Recha et al. 2012) and catchments monitored continuously. The boundaries of each catchment were determined using a Global Positioning System (GPS) on the landscape around each spring up to the plateau, using an average of five to eight independent assessments. The GPS data were then used to generate a Geographical Information System (GIS) output and map of the area. The sizes of the catchments were 12.8 ha for the forest, 14.4 ha for the 5-year old conversion, 9.1 ha for the 10 year conversion and 10.0 ha for the 50-year conversion. There were a total of one, six, and eleven households living in the 5-, 10-, and 50-year conversion catchments, respectively. A standard V-notch weir was constructed at each catchment outlet for determining stream discharge.

Stream stage was recorded using water capacitance probes (Odyssey Dataflow Systems Pty Ltd., New Zealand) installed at the weir. The probes were programmed to give a reading of the average stream stage at 2 min. Data from these probes were downloaded biweekly. The weir ratings were determined at low and intermediate flows. The weir rating correlation coefficients were $r^2 = 0.944$, $r^2 = 0.915$, $r^2 = 0.926$, $r^2 = 0.931$ for the forest, 5-, 10- and 50-year conversions, respectively. The stream hydrographs were normalized by corresponding catchment sizes to allow comparison between responses for the 4 catchments.

Stream water sampling was done biweekly at the weir outlet, at the beginning and mid of every month. The water samples were filtered through 0.45- μ m pore-size glass-fiber filter, into two separate 50-mL centrifuge vials. We added thymol into the first 50-mL centrifuge vial that was used for determination of calcium (Ca), magnesium (Mg), sodium (Na), potassium (K), total dissolved phosphorus (TDP), nitrate nitrogen (NO₃-N), ammonium nitrogen (NH₄⁺-N) and total dissolved nitrogen (TDN). We added 10 % HCl into the second 50-mL centrifuge vial that was used for determination of dissolved organic carbon (DOC) and the DOC stable isotopic ratio of ¹³C to ¹²C. The 0.45- μ m pore-size glass-fiber filter paper with sediment was air dried and kept for the determination of coarse particulate organic carbon (CPOC), the CPOC isotopic ratio of ¹³C to ¹²C, and coarse particulate nitrogen (CPON).

Five soil samples from 0 to 0.1 m depth were collected randomly by stratifying for location within slope and plateau of each catchment, after carefully removing the litter layer (capturing only the mineral horizon). They were analyzed for SOM, SOC, total N, available P, Ca, K, Mg and Na. Similarly, four sites were randomly selected by stratifying for location within slope and plateau of each catchment and additional soil sampled at depths of 0.1–0.3, 0.3–0.9, 0.9–1.5, 1.5–2.4 m. Bulk density was determined by the soil core method; we used five soil samples from a depth of 0–0.1 m in each catchment (Campbell and Henshall 1991). Soil porosity (ϕ) was computed from bulk density (ρ_b) and particle density (ρ_p) using the formula $\phi = 1 - (\rho_b/\rho_p)$. The use of organic and inorganic fertilizers was assessed in all studied agricultural watersheds by a full survey of all households using interviews. Farmer interviews in the watersheds confirmed information provided by the studies of Kinyangi (2008), Kimetu (2009),

Fig. 1 Map of the study area of the Kapchorwa headwater chronosequence of catchments in western Kenya, and its approximate location within Kenya (inset shows Kenya and its districts). The weir positions are shown by dots in each catchment. Letters indicate land use histories of the catchments: forested catchment (a), maize cultivation for 5 years (b), 10 years (c) and 50 years (d) following forest conversion



and Moebius-Clune et al. (2011). The soils received nil, or only very little inorganic fertilizer since forest clearing, no animal manure, and were always cropped with maize. From the interviews, only di-ammonium phosphate (DAP, N:P:K of 18:46:0) fertilizer was used at an application rate of 6 and 9 kg ha⁻¹ within the 10 year and 50 year watersheds, respectively, for the year 2008. The farmers in the 5 year watershed did not use any fertilizer. The N and P addition was 1.02 and 2.76 kg ha⁻¹, respectively, for the 10 year watershed. The addition for the 50 year watershed was 1.62 and 4.14 kg ha⁻¹ of N and P, respectively. Animal manure was not used in any household.

Laboratory measurements

Soil total C was determined by dry combustion after fine grinding soil using a Mixer Mill (MM301, Retsch, Germany). Samples were analyzed for total C contents with a Europa ANCA-GSL CN analyzer (PDZ Europa Ltd., Sandbach, UK). Soil organic matter (SOM) was determined by loss on ignition (Storer 1984) and soil pH (in water) at the w/v ratio 1:2.5 using a glass electrode (Thermo Scientific, Beverly, MA, US). Filters containing CPOC were ground using a Mixer Mill (MM301, Retsch, Germany) and analyzed for total C, the isotopic ratio of ¹³C to ¹²C, and CPON using a coupled Europa 20–20 continuous isotope ratio mass spectrometry (PDZ Europa Ltd., Sandbach, UK). The Mehlich 3 extraction procedure (Mehlich 1984) was used for the available soil P, Ca, Mg, K and Na. The Ca, Mg, K, Na and P concentrations were obtained by Inductively Coupled

Plasma Atomic Emission Spectroscopy (ICP-AES, ARCOS, Germany). TDN was analyzed using Shimadzu's Total Nitrogen Module, TNM-1 that uses chemiluminescence. Dissolved organic nitrogen (DON) was computed using the formula $DON = TDN - NO_3^- + NH_4^+$. DOC analysis was carried out on a Shimadzu Total Organic Carbon-Visionary Series (TOC-V_{CSH}) analyzer following the procedure described by Qian and Mopper (1996). NO₃-N and NH₄⁺-N were determined on a Seal AQ2-Automated Discreet Analyzer (Seal Analytical, England).

Statistical analyses

Statistical analyses were done using JMP Version 8 (SAS Institute Inc, Cary, NC, USA) for soil properties and stream water solutes of the four catchments. A one way ANOVA was used for the soil properties, and a repeated measure analysis controlling for date of sampling was performed for the stream water solutes. Both analyses were followed with posthoc multiple comparisons using a Tukey correction for multiple comparisons when the catchment effect was significant. Comparisons were considered significant at $P < 0.05$.

Results

Soil properties

Within the top 0.1 m, the SOM and SOC concentrations significantly decreased by 59 % and 75 %, respectively.

Table 1 Soil properties in the top 0.1 m

	Forest	5 year conversion	10 year conversion	50 year conversion
Soil organic matter (%)	17.08a	13.67b	8.53c	7.00d
Soil organic carbon (g kg ⁻¹)	108.3a	68.8b	36.4c	27.5c
Soil organic carbon (t ha ⁻¹)	86.6a	62.6b	37.5c	32.2c
Soil nitrogen (g kg ⁻¹)	11.2a	6.9b	3.4c	2.3c
Soil nitrogen (t ha ⁻¹)	8.96a	6.28b	3.50c	2.69c
Soil bulk density (g cm ⁻³)	0.80c	0.91b	1.03ab	1.17a
Total porosity	0.70a	0.66b	0.61bc	0.56c
Moisture at 0.33 bar (% v/v, field capacity)	34.7a	34.1a	24.9b	23.0b
pH (water, 1:2.5)	7.39a	6.48b	6.23b	5.81c
Available P (g kg ⁻¹)	0.011a	0.007a	0.003a	0.006a
Available K (g kg ⁻¹)	0.69a	0.58ab	0.23b	0.44ab
Available Ca (g kg ⁻¹)	6.25a	4.98a	1.77b	2.91b
Available Mg (g kg ⁻¹)	0.73a	0.45b	0.20c	0.33bc
Available Na (g kg ⁻¹)	0.026a	0.020a	0.010a	0.011a

Means within a row followed by the same letters are not significantly different from each other at $P < 0.05$ ($n = 5$)

respectively, and the soil N contents by 79 % from the forest to the 50-year old agricultural catchment (Table 1). The forest SOC and soil N were significantly ($P < 0.05$) higher than the 5-year conversion. Soils in the 10- and 50-year conversion did not differ. The SOM, SOC and N loss was very rapid in the first 10 years of conversion decreasing by 50, 66 and 70 %, respectively. Available soil P, K and Na did not change as a result of forest clearing and soil use between the studied watersheds. In contrast, extractable Ca and Mg significantly ($P < 0.05$) decreased over time (Table 1). Similarly, pH values significantly decreased from forest (7.39) to agriculture (5.81).

The soil ρ_b increased rapidly by 28.8 % in the first 10 years of cultivation from 0.80 to 1.03 g cm⁻³ (Table 1). Overall, the soil ρ_b increased by 46.3 % from 0.8 to 1.17 within 50 years of cultivation. The rapid increase in ρ_b in the first 10 years after forest conversion was followed by a 12 % drop in the total porosity from 0.69 to 0.61 in the same period.

In the 0–0.1 m depth, the soil $\delta^{13}\text{C}$ values were -26.07 ± 0.68 ‰, -25.37 ± 0.84 ‰, -21.21 ± 0.87 ‰ and -19.18 ± 0.44 ‰ for the forest, 5-, 10- and 50-year conversion catchments, respectively (Fig. 2). This indicates enrichment in the heavier isotope from the forest to the catchment under cultivation for 50 years. The $\delta^{13}\text{C}$ values at the topsoil (0–0.1 m) increased linearly over time from -26.07 to -19.18 ‰ ($r^2 = 0.95$; $P < 0.0001$), with similar

trends at depth. Similarly, the $\delta^{13}\text{C}$ values became less negative deeper in the profile for all the catchments indicating a slight enrichment in the heavier isotope compared to the topsoil.

Stream water chemistry

The $\delta^{13}\text{C}$ composition of CPOC (Table 2) did not change from the forest (-27.52 ‰) to the 10 year conversion (-27.28 ‰). The 50-year old catchment had a slightly higher $\delta^{13}\text{C}$ value compared to the other catchments. The streamwater DOC $\delta^{13}\text{C}$ composition




Fig. 2 Long term shift in the isotopic composition of $\delta^{13}\text{C}$ of the soils at various depths in the watersheds. $n = 4$. Bars are ± 1 standard deviation of the mean for each depth

Table 2 The isotopic composition of $\delta^{13}\text{C}$ of the CPOC and DOC, and the flow weighted nutrient concentrations of the headwater catchments

	Forest	5 year conversion	10 year conversion	50 year conversion
<i>Sediment and filtered streamwater</i>				
CPOC $\delta^{13}\text{C}$ (‰)	-27.52b	-27.29b	-27.28b	-26.40a
DOC $\delta^{13}\text{C}$ (‰)	-28.59c	-28.25c	-26.79b	-22.75a
<i>Streamflow concentrations</i>				
DOC (mg L^{-1})	1.31a	1.48a	1.47a	1.52a
CPOC (mg L^{-1})	1.25a	1.43a	1.49a	1.65a
CPON (mg L^{-1})	0.12a	0.14a	0.15a	0.16a
TDN (mg L^{-1})	0.49b	0.67b	4.83a	4.64a
NO_3^- (mg L^{-1})	0.40b	0.58b	4.71a	4.52a
DON (mg L^{-1})	0.10a	0.09a	0.10a	0.12a
TDP (mg L^{-1})	0.03b	0.01b	0.05b	0.15a
K (mg L^{-1})	0.36c	0.80b	0.95a	1.18a
Ca (mg L^{-1})	7.23a	5.55b	5.48b	7.16a
Mg (mg L^{-1})	3.01a	2.04c	1.61d	2.54b
Na (mg L^{-1})	2.99c	3.70c	4.65b	5.61a

Means within a row followed by the same letters are not significantly different from each other at $P < 0.05$ ($n = 24$ for CPOC $\delta^{13}\text{C}$, $n = 4$ for DOC $\delta^{13}\text{C}$, and $n = 12$ for all others)

followed the same trend as the sediment but with a significantly greater magnitude of enrichment in the heavier isotope from the forest and recent conversion to the 10-year old conversion. Discharge in the 50-year conversion had a significantly higher value than that of all other headwaters (-22.75 ‰). The flow-weighted average concentrations of DOC, CPOC and CPON did not change with forest conversion and duration of cultivation (Table 2). TDN concentrations in the discharge from forest and 5-year catchments did not differ, but increased with longer cultivation. NO_3^- was the dominant form of dissolved N (91 %) in the Kapchorwa watershed fluvial ecosystem, followed by DON (9 %), whereas NH_4^+ was below our detection limit. The 10- and 50-year conversions had nine times higher TDN concentrations compared to the forest and most recent conversion. Stream TDP concentrations were at least one order of magnitude lower compared to TDN, Ca, K, and Mg. TDP concentrations of the forest, 5- and 10-year conversions were threefold lower than that of the 50-year conversion ($P < 0.05$). Stream water K and Na concentrations both increased upon forest conversion and subsequent cultivation. Overall, K concentrations increased threefold and Na concentrations twofold following 50 years of cultivation after forest conversion. In contrast to all other solutes, Ca and Mg concentrations did not show a consistent trend from forest to cultivation.

The total C exports increased after forest conversion and with subsequent cultivation (Table 3). The DOC and CPOC export doubled in the first 5 years following forest conversion. The increase continued by a further 43 and 70 % from the 5- to the 50-year conversions, respectively. Amounts of CPON, TDN and NO_3^- exported were also twice the first 5 years after forest conversion. Upon longer cultivation between 5 and 10 years, CPON export increased 3.5 times, while the TDN and NO_3^- export increased eightfold. Between 10 and 50 years of continuous cultivation, the amount of CPON export doubled while that of TDN and NO_3^- did not increase further. The magnitude of TDP export was far less than that of C and all other nutrients. The amounts of TDP exported did not change after forest conversion ($<0.03 \text{ kg ha}^{-1}$), but increased with subsequent cultivation up to the 0.98 kg ha^{-1} . The cations (K, Ca Mg and Na) exhibited a similar trend whereby the amounts exported increased from the forest to the 5-, 10- and 50-year conversions. Among the measured base cations, K had the lowest export while Ca had the highest.

There was not a clear separation of the hydrograph between the long rainy (April-June) and short rainy (August-October) seasons in the Kapchorwa catchment in 2008. The stream flow response to precipitation events increased with longer cultivation (Fig. 3). There were no relevant changes in the concentrations of DOC, TDN, TDP, Ca, and Mg across the whole

Table 3 Streamwater carbon and nutrient exports ($\text{kg ha}^{-1} \text{ year}^{-1}$) from the Kapchorwa headwater catchments

	Forest	5 year conversion	10 year conversion	50 year conversion
DOC	3.87	6.85	8.46	9.79
CPOC	3.94	6.60	8.57	10.63
CPON	0.33	0.64	0.87	1.03
TDN	1.45	3.12	27.76	29.93
NO_3^-	1.18	2.68	27.09	29.16
DON	0.27	0.44	0.67	0.77
TDP	0.09	0.03	0.29	0.98
K	1.06	3.68	5.44	7.61
Ca	21.4	25.7	31.5	46.2
Mg	8.93	9.42	9.23	16.38
Na	8.86	17.08	26.70	36.22

year. The K and Na concentrations, however, decreased slightly after the long rainy season in August and remained low for the short rainy season. Overall, there was no relationship between the solute concentrations and discharge (Fig. 4).

Discussion

Sources of fluvial OC and nutrient losses

Even though annual fluvial OC exports increased more than three-fold (our study) and CO_2 evasion from SOC mineralization decreased (Kimetu and Lehmann 2010), annual DOC and CPOC losses by stream water of 22 kg ha^{-1} from the highly degraded soils were still low with only 2 % mineralization losses of the $1,200 \text{ kg ha}^{-1}$ determined by direct measurements from the soil surface using static vented chambers. Similarly, cumulative fluvial organic C losses were 0.9 t ha^{-1} (2 %) over the first 50 years (using linear interpolation of the sum of DOC and CPOC) compared to SOC losses of 54 t ha^{-1} in the first 0.1 m (calculated from Table 1; similar SOC losses reported by Solomon et al. 2007; Kimetu and Lehmann 2010). It is possible that inorganic C and particularly dissolved CO_2 may significantly contribute to total fluvial C losses from watersheds as shown in the Amazon (Johnson et al. 2008). However, even with a possibly larger inorganic than organic C loss, microbial mineralization from the soil still exceeds 90 % of total SOC losses over the 50 years. The quantified

fluvial C export is low compared to other studies. Moore et al. (2010) observed a fluvial total organic C removal of 12 % by a Bornean blackwater river in a peat-covered area of Indonesia. Cumulative fluvial N losses over the first 50 years (using linear interpolation of the sum of TDN and CPON) were 1.3 t ha^{-1} compared to the total soil N losses of 6.3 t ha^{-1} (representing 21 %) in the first 0.1 m (calculated from Table 1). Similar to fluvial C losses, watersheds at our site showed lower export than at other locations.

Both fluvial CPOC and DOC were mainly mobilized from the topsoil, as $\delta^{13}\text{C}$ values in stream discharge were lower than in the topsoil and soil values increased with depth as also reported by Krull et al. (2002) at a nearby site. Similar conclusions were drawn by Frank et al. (2000) and Raymond and Saiers (2010) who found most of the DOC losses to be derived from the topsoil in catchments in central Switzerland and eastern United States, respectively. Alternatively, the low $\delta^{13}\text{C}$ values in our study may also indicate that OC losses originated from older SOC fractions characterized by small size and association with minerals that showed lower $\delta^{13}\text{C}$ values in the same chronosequence (Kinyangi 2008). The isotopic composition of the topsoil correlated strongly with the DOC $\delta^{13}\text{C}$ ($r^2 = 0.84$; $y = 1.132x + 7.1478$) as well as with the CPOC $\delta^{13}\text{C}$ ($r^2 = 0.83$; $y = 18.25x - 49.523$), but with a low slope for CPOC.

Nutrient losses, crop production and stream pollution

Total N (TDN + CPON) losses of $31 \text{ kg ha}^{-1} \text{ year}^{-1}$ in the highly degraded soils were relevant compared to crop N uptake of maize at the same sites with $90 \text{ kg N ha}^{-1} \text{ year}^{-1}$ (Kimetu et al. 2008). There was no fertilizer application in the recent conversion, while applications in the 10- and 50-year conversions were about 40 kg N ha^{-1} (average from all farms in the studied catchments). The typical application rates to maize by most farmers in the entire region are 20 kg N ha^{-1} (Ngoze 2008), less than the stream water losses reported here. Even more dramatic were the fluvial Ca and Mg losses of 46 and $16 \text{ kg ha}^{-1} \text{ year}^{-1}$, respectively, which were greater than uptake by a single maize crop of 20 and 10 kg ha^{-1} , respectively (Kimetu et al. 2008). In contrast, annual fluvial P losses of 1 kg ha^{-1} were low compared to plant uptake of 10 kg ha^{-1} (Kimetu et al. 2008) and




Fig. 3 Bi-weekly discharge and concentrations of DOC, Na, Ca, K and Mg of the Kapchorwa headwater catchments




Fig. 4 The relationship between discharge and DOC, Na and nutrient concentrations in the 50 year watershed

fertilization of about 7.5 kg P ha⁻¹ (sum of inorganic and organic fertilizers averaged for all farms) in the oldest conversion. However, redistribution of P within

the watershed was not measured and stream water losses are only a lower boundary of actual losses of soil P. Nonetheless, both Ca and Mg as well as P losses by stream water may not be as serious as N losses because available P, Ca, and Mg in soil decreased much less than total N (Table 1; similar for the same sites reported by Ngoze et al. 2008; Kinyangi 2008), and crops were found to respond less to P than to N additions (Ngoze et al. 2008). In addition, TDN concentrations and total losses by stream discharge increased the most by forest conversion and continuous cultivation compared to any other nutrient studied here.

Despite the significant N losses from stream discharge for agricultural productivity, the observed concentrations in the studied headwater catchments are not an environmental concern for aquatic environments. Average annual N concentrations at or below 3 mg L⁻¹ and maximum concentrations of 6 mg L⁻¹ during the dry season of December to March (Fig. 2) do not pose a risk for human health

(Rubio-Arias et al. 2010) and are not expected to cause eutrophication (Hill et al. 2011; King and Balogh 2011). Similarly, P concentrations of 0.01–0.15 mg P L⁻¹ (Fig. 2) in the stream discharge are moderate but not a cause for significant environmental concern (Auer et al. 1986; King and Balogh 2011; Yu et al. 2011). Values up to 6.7 mg PO₄-P L⁻¹ have been found in the River Thames catchment in the UK (Young et al. 1999) and up to 10 mg PO₄-P L⁻¹ in the Spree River in Germany (Lewandowski and Nutzmann 2010). Also DOC concentrations are significantly lower in the studied headwaters than in the lower Yala and Lake Victoria (Lalah and Wandiga 2006). Therefore, the well documented contamination of the lower Yala River (Aloo 2003) and Lake Victoria with DOC, N and P which has been suspected to cause the observed eutrophication (Scheren et al. 2000; Lalah and Wandiga 2006; Hecky et al. 2010) does not seem to be caused by agriculture in the headwater catchments, but must occur further downstream.

Carbon and nutrient losses after landuse change

Our DOC concentrations and total losses were in the same range as reported by Lesack et al. (1984) from the Gambia River in the West African savannah with a DOC concentration of 1.98 mg L⁻¹ and DOC export of 2.67 kg ha⁻¹ year⁻¹. Similar DOC concentrations but 2–10 times greater export were found by Cairns et al. (2009) in the forests of the Oregon Cascades and of southern Amazonia by Johnson et al. (2006b), yet values are within the range reported for a variety of watersheds elsewhere (Lesack et al. 1984; Selva et al. 2007; Stutter et al. 2008; Waldron et al. 2009). The CPOC and CPON concentrations and export were in the same order of magnitude as those of DOC and DON. This finding differs from that of Vidal-Abarca et al. (2001) who found DOC to be the most important fraction (98 %) of organic C flowing in a saline and semi-arid stream in Spain. Similarly greater proportions of DOC were also reported by Johnson et al. (2006a, b) from southern Amazonia. Therefore, particulate OC losses are greater than in other studies, but the observed fluvial OC losses from the forest seem to be representative of those found in other studies. The examined change of solute losses as a response to landuse change may therefore be indicative of losses in other regions.

The increase in streamwater concentrations of TDN, NO₃⁻ and DON with longer duration of cultivation can be attributed to three sources. (1) Mineralization of SOM led to accumulation of mineral N and especially nitrate (as seen from stream water concentrations) which is weakly adsorbed to soil minerals and easily leached. (2) The greater number of households on older conversions presumably led to higher inputs of farm manure from the cattle kept by the farmers. The manure was shown to consist of 22 g N kg⁻¹ (Kimetu et al. 2008). Half of the farmers in the older conversions applied inorganic fertilizer equivalent to 40 kg N ha⁻¹. (3) As stated above, only farmers with fields that have been cultivated for 10 years or longer apply mineral fertilizers. Fields that have been recently cleared at the same sites are rarely fertilized due to their high N mineralization rates and their low response to mineral (Ngoze et al. 2008) and organic N applications (Kimetu et al. 2008).

The increase in fluvial K and Na concentrations with longer cultivation may be explained by lower SOC contents and the soil mineralogy being dominated by kaolinite, both resulting in low cation retention (Kimetu and Lehmann 2010). Krull et al. (2002) reported mineralogy dominated by quartz, kaolin and mica, with minor components associated with feldspars and oxides in the adjacent Kakamega part of the forest. They observed an abundance of quartz, kaolinite, muscovite and microcline with a slight increase in goethite with increasing soil depth. The observed greater proportion of surface runoff of 4–10 % of rainfall from forest compared to long-term cultivated fields (Recha et al. 2012) may also have led to a greater mobilization of K and Na near the surface. Extractable K concentrations did not decrease with forest conversion and cultivation (Table 1; compare with Kinyangi 2008). Similar to N, a greater number of households in the older conversion watersheds may have added more kitchen waste that are rich in K and Na such as ash from firewood cookstoves and farm yard manure. Kimetu et al. (2008) documented contents of 23.2 g K kg⁻¹ in the manure.

The lack of change or even slight decrease in Ca and Mg stream water concentrations may be a result of a dominance of weathering as the primary source of Ca and, to a certain extent, Mg. In addition, the significant decrease in extractable Ca in the topsoil (Table 1) in combination with the rather decreasing stream water

concentrations after forest conversion point at the subsoil as the primary source of Ca.

Intra-annual dynamics of stream water solutes

None of the solute concentrations (except DOC, Na and K) changed with the strong changes in discharge between January and December 2008 (Figs. 3 and 4). No significant correlations were established between any of the solutes and stream discharge (Fig. 4) which is in contrast to other studies that typically show significant changes in DOC and nutrient concentrations throughout the year (Markewitz et al. 2001; Johnson et al. 2006b; Bucker et al. 2011). Although variations in intra-annual stream concentrations are often observed, the solutes can either be more concentrated during high flow in the rainy season due to export of mainly surficial nutrients (Markewitz et al. 2001) or less concentrated due to dilution during heavy rainfall and larger volume of stream discharge (Elsenbeer et al. 1994; Anderson et al. 1997; Tsujimura et al. 2001; Grimaldi et al. 2004). The constant concentration of N, P, Mg and Ca with varying discharge over seasonal time scales in our study may be explained by a balance between changes in water delivery and solute delivery (Salmon et al. 2001). Ca and Mg can be delivered from both organic-rich soil horizons and weathering of deeper soil (Yusop et al. 2006). During low flows, the Ca and Mg would be derived from the weathered mineral subsoil, whereas near surface storm event flow (Noguchi et al. 1997) could entrain these solutes from the organic-rich topsoil.

In contrast, the Na and K concentrations in all four catchments dropped during the month of August at the onset of the second rainy season. The sources of K as well as Na are associated with leaching and decomposition of leaves and organic matter, and thus are expected to be available at greater quantities in the upper soil profile (Proctor et al. 1989; Veneklaas 1991). Therefore, prolonged rain may have leached K and Na contained in the litter layer from the soil surface similar to what has been observed for DOC in southern Amazonia (Johnson et al. 2006b) where the litter layer disappeared towards the end of the rainy season (Selva et al. 2007).

Conclusion

Results from this study indicate that conversion of forest catchments to continuous maize cropping had a

significant effect on the hydrochemistry of headwater streams. All OC species and nutrient concentrations increased with forest conversion and cultivation except for Ca and Mg. The reason may be the proportional greater importance of geochemical weathering for Ca and Mg. Dilution of major solutes during the rainy season frequently observed elsewhere did not occur here, except for K and Na. Fluvial OC export was low compared to estimated mineralization of SOC and stream P losses were negligible, but N losses by stream water must be addressed to curtail declining crop productivity. In terms of nutrient pollution and possible contamination of water resources, the N and P levels are not of concern and agriculture in the headwaters seems to play a minor role for the observed eutrophication of the downstream Yala River system and Lake Victoria.

Acknowledgments The authors would like to thank Brett Gleitsmann for assistance in identifying the study sites, constructing the weirs and doing initial sampling. We thank Henry Biwott for help with field work and Stephen DeGloria for help with GIS. This research was financially supported in part by grants from the Ford Foundation International Fellowships Program, the National Science Foundation Grant No. 0215890 through CIFAD, the Biogeochemistry and Environmental Biocomplexity Small Grant Program Ref. DGE 0221658, Towards Sustainability Fund, the Institute for African Development and the Norman Borlaug Leadership Enhancement in Agriculture Program Fellowship. Logistical support was provided by the World Agroforestry Centre Kisumu office. We thank two anonymous referees for their valuable suggestions.

References

- Aloo PA (2003) Biological diversity of the Yala Swamp lakes, with special emphasis on fish species composition, in relation to changes in the Lake Victoria Basin (Kenya): threats and conservation measures. *Biodivers Conserv* 12:905–920
- Anderson SP, Dietrich WE, Torres R, Montgomery DR, Loague K (1997) Concentration-discharge relationships in runoff from a steep, unchanneled catchment. *Water Resour Res* 33:211–225
- Auer MT, Kieser MS, Canale RP (1986) Identification of critical nutrient levels through field verification of models for phosphorus and phytoplankton growth. *Can J Fish Aquat Sci* 43:379–388
- Awiti AO, Walsh MG, Kinyamario J (2008) Dynamics of tropical carbon and nitrogen along a tropical forest-cropland chronosequence: evidence from stable isotope analysis and spectroscopy. *Agric Ecosyst Environ* 127:265–272
- Bleher B, Uster D, Bergsdorf T (2006) Assessment of threat status and management effectiveness in Kakamega forest, Kenya. *Biodivers Conserv* 15:1159–1177

- Bucker A, Crespo P, Frede H, Breuer L (2011) Solute behavior and export rates in neotropical montane catchments under different land-uses. *J Trop Ecol* 27:305–317
- Cairns MA, Lajtha K, Beedlow PA (2009) Dissolved carbon and nitrogen losses from forests of the Oregon Cascades over a successional gradient. *Plant Soil* 318:185–196
- Campbell DJ, Henshall JK (1991) Bulk density. In: Smith KA, Mullins CE (eds) *Soil analysis-physical methods*. Marcel Dekker, New York, pp 329–366
- Carpenter SR, Caraco NF, Correll DL, Howarth RW, Sharpley AN, Smith VH (1998) Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecol Appl* 8:559–568
- Davidson EA, Ackerman IL (1993) Changes in soil carbon inventories following cultivation of previously untilled soils. *Biogeochemistry* 20:161–193
- Davidson EA, Neill C, Krusche AV, Ballester VVR, Markewitz D, de Figueiredo RO (2004) Loss of nutrients from terrestrial ecosystems to streams and atmosphere following land use change in Amazonia. In: DeFries R, Asner G, Houghton R (eds) *Ecosystems and land use change*. Geophysical Monographs, AGU, Washington, pp 147–158
- deGraffenried JB, Jr Shepherd KD (2009) Rapid erosion modeling in a western Kenya watershed using near infrared reflectance, classification tree analysis and ¹³⁷Cesium. *Geoderma* 154:93–100
- Elsenbeer H (2001) Hydrologic flowpaths in tropical rainforest landscapes—a review. *Hydrol Process* 15:1751–1759
- Elsenbeer H, West A, Bonell M (1994) Hydrologic pathways and stormflow hydrochemistry at South Creek, Northeast Queensland. *J Hydrol* 162:1–21
- FAO-UNESCO-ISRIC (1988) Revised legend, FAO-UNESCO Soil map of the world. *World Soil Resources Report* 60, Rome, 119 pp
- Frank H, Patrick S, Peter W, Hannes F (2000) Export of dissolved organic carbon and nitrogen from Gleysol dominated catchments—the significance of water flow paths. *Biogeochemistry* 50:137–161
- Germer S, Neill C, Krusche AV, Elsenbeer H (2010) Influence of land-use change on near-surface hydrological processes: undisturbed forest to pasture. *J Hydrol* 380:473–480
- Glenday J (2006) Carbon storage and emissions offset potential in an East African tropical rainforest. *For Ecol Manag* 235:72–83
- Grimaldi C, Drimaldi M, Millet A, Bariac T, Boulegue J (2004) Behaviour of chemical solutes during a storm in a rainforested headwater catchment. *Hydrol Process* 18:93–106
- Grip H, Fritsch JM, Bruijnzeel LA (2004) Soil and water impacts during forest conversion and stabilization to new land use. In: Bonell M, Bruijnzeel LA (eds) *Forests, water, and people in the humid tropics: past, present, and future*. Hydrological Research for Integrated Land and Water Management, UNESCO International 531 Hydrology Series, Cambridge University Press, Cambridge, UK, pp 561–589
- Hartemink AE, Veldkamp T, Bai ZG (2008) Land cover change and soil fertility decline in tropical regions. *Turkish J Agric For* 32:195–213
- Hecky RE, Mugidde R, Ramlal PS, Talbot MR, Kling GW (2010) Multiple stressors cause rapid ecosystem change in Lake Victoria. *Freshw Biol* 55(Suppl 1):19–42
- Hertel D, Moser G, Culmsee H, Erasmi S, Horna V, Schuldt B, Leuschner C (2009) Below and above ground biomass and net primary production in a paleotropical natural forest (Sulawesi, Indonesia) as compared to neotropical forest. *For Ecol Manag* 258(9):1904–1912
- Hill BH, Bolgrien DW, Herlihy AT, Jicha TM, Angradi TR (2011) A synoptic survey of nitrogen and phosphorus in tributary streams and great rivers of the Upper Mississippi, Missouri, and Ohio river basins. *Water Air Soil Pollut* 216:605–619
- Huggett RJ (1998) Soil chronosequences, soil development, and soil evolution: a critical review. *Catena* 32:155–172
- Jaetzold R, Schmidt H (1983) *Farm management handbook of Kenya, vol 2 Natural Conditions and Farm Management Information, Part A*. West Kenya. Ministry of Agriculture and Livestock Development, Kenya, 319 pp
- Johnson MS, Lehmann J, Couto EG, Novaes-Filho JP, Riha S (2006a) DOC and DIC in flowpaths of Amazonian headwater catchments with hydrologically contrasting soils. *Biogeochemistry* 81:45–57
- Johnson MS, Lehmann J, Selva EC, Abdo M, Riha S, Cuoto EG (2006b) Organic carbon fluxes within and streamwater exports from headwater catchments in the southern Amazon. *Hydrol Process* 20:2599–2614
- Johnson MS, Lehmann J, Riha SJ, Krusche AV, Richey JE, Ometto JPHB, Couto EG (2008) CO₂ efflux from Amazonian headwater streams represents a significant fate for deep soil respiration. *Geophys Res Lett* 35:L17401
- Kenya National Bureau of Statistics (2010) *The Kenya census 2009 report*. Government of Kenya. <http://www.knbs.or.ke/>. Accessed 15 May 2012
- Kimetu JM (2009) *Soil organic matter revitalization: implications on food production and security in tropical agricultural systems*. PhD dissertation, Cornell University, Ithaca, NY, USA
- Kimetu JM, Lehmann J (2010) Stability and stabilisation of biochar and green manure in soil with different organic carbon contents. *Aust J Soil Res* 48:577–585
- Kimetu JM, Lehmann J, Ngoze S, Mugendi D, Kinyangi J, Riha S, Verchot L, Recha JW, Pell A (2008) Reversibility of soil productivity decline with organic matter of differing quality along a degradation gradient. *Ecosystems* 11(5):726–739
- Kimetu JM, Lehmann J, Kinyangi JM, Cheng CH, Thies J, Mugendi DN, Pell A (2009) Soil organic C stabilization and thresholds in C saturation. *Soil Biol Biochem* 41:2100–2104
- King KW, Balogh JC (2011) Stream water nutrient enrichment in a mixed-use watershed. *J Environ Monit* 13:721–731
- Kinyangi JM (2008) *Soil degradation, thresholds and dynamics of long-term cultivation: from landscape biogeochemistry to nanoscale biogeocomplexity*. PhD dissertation Cornell University
- Krull ES, Bestland EA, Gates WP (2002) Soil organic matter decomposition and turnover in a tropical ultisol: evidence from $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and geochemistry. *Radiocarbon* 44(1):93–112
- Lal R (2006) Enhancing crop yields in the developing countries through restoration of the soil organic carbon pool in agricultural lands. *Land Degrad Dev* 17:197–209
- Lalah JO, Wandiga SO (2006) Extinction coefficients and dissolved organic carbon content in freshwater in Kenya. *Bull Environ Contam Toxicol* 77:533–542

- Lesack LFW (1993) Export of nutrients and major ionic solutes from a rain forest catchment in central Amazonian Basin. *Water Resour Res* 29(3):743–758
- Lesack LFW, Hecky RE, Melack JM (1984) Transport of carbon, nitrogen, phosphorus, and major solutes in the Gambia River, West Africa. *Limnol Oceanogr* 29:816–830
- Lewandowski J, Nutzman G (2010) Nutrient retention and release in a floodplain's aquifer and in the hyporheic zone of a lowland river. *Ecol Eng* 36:1156–1166
- Likens GE, Bormann FH (1995) *Biogeochemistry of a forested ecosystem*, 2nd edn. Springer, New York 159p
- Lung T, Schaab G (2006) Assessing fragmentation and disturbance of west Kenyan rainforests by means of remotely sensed time series data and landscape metrics. *Afr J Ecol* 44:491–506
- Malmer A (1996) Hydrological effects and nutrient losses of forest plantation establishment on tropical rainforest land in Sabah, Malaysia. *J Hydrol* 174:129–148
- Malmer A, Grip H (1990) Soil disturbance and loss of infiltrability caused by mechanized and manual extraction of tropical rainforest in Sabah, Malaysia. *For Ecol Manag* 38:1–12
- Mann LK (1986) Changes in soil carbon after cultivation. *Soil Sci* 142:279–288
- Markewitz D, Davidson EA, Figueiredo RO, Victoria RL, Krusche AV (2001) Control of cation concentrations in stream waters by surface soil processes in an Amazonian watershed. *Nature* 410:802–805
- Markewitz D, Davidson EA, Moutinho P, Nepstad D (2004) Nutrient loss and redistribution after forest clearing on a highly weathered soil in Amazonia. *Ecol Appl* 14(Suppl 4):S177–S199
- McLaughlan K (2006) The nature and longevity of agricultural impacts on soil carbon and nutrients: a review. *Ecosystems* 9:1364–1382
- Mehlich A (1984) Mehlich 3 soil test extract: a modification of Mehlich 2 extractant. *Comm Soil Sci Plant Anal* 15:1409–1416
- Moebius-Clune BN, van Es HM, Idowu OJ, Schindelbeck RR, Kimetu JM, Ngoze S, Lehmann J, Kinyangi JM (2011) Long-term soil quality degradation along a cultivation chronosequence in Western Kenya. *Agric Ecosyst Environ* 141:86–99
- Moore S, Gauci V, Evans CD, Paage SE (2010) Fluvial organic carbon losses from a Bornean blackwater river. *Biogeosci Discuss* 7:8319–8343
- Nair VD, Graetz DA (2004) Agroforestry as an approach to minimizing nutrient loss from heavily fertilized soils: the Florida experience. *Agrofor Syst* 61:269–279
- Ngoze SO (2008) Soil nutrient depletion and repletion in a tropical agroecosystem. PhD dissertation, Cornell University
- Ngoze SO, Riha S, Lehmann J, Verchot L, Kinyangi J, Mbugua D, Pell A (2008) Nutrient constraints to tropical productivity in long-term degrading soils. *Global Change Biol* 14:1–13
- Noguchi S, Abdul Rahim N, Baharuddin K, Sammori T, Tani M, Morisada S (1997) Rainfall-runoff responses and role of soil moisture variations to the response in tropical rain forest, Bukit Tarek Peninsular Malaysia. *J For Resour* 2:115–120
- Pandey CB, Singh GB, Singh SK, Singh RK (2010) Soil nitrogen and microbial biomass carbon dynamics in native forests and derived agricultural land uses in humid tropical climate of India. *Plant Soil* 333:453–467
- Proctor J, Phillips C, Duff GK, Heaney A, Robertson FM (1989) Ecological study of Gunung Silam, a small ultrabasic mountain in Sabah Malaysia II. Some forest processes. *J Ecol* 77:317–331
- Qian J, Mopper K (1996) An automated, high performance, high temperature combustion dissolved organic carbon analyzer. *Anal Chem* 68(18):3090–3097
- Raymond PA, Saiers JE (2010) Event controlled DOC export from forested watersheds. *Biogeochemistry* 100:197–209
- Recha J, Lehmann J, Walter M, Pell A, Verchot L, Johnson M (2012) Stream discharge in tropical headwater catchments as a result of forest clearing and soil degradation. *Earth Interact*: published online. doi:10.1175/2012EI000439.1
- Richey JE, Melack JM, Aufdenkampe AK, Ballester VM, Hess LL (2002) Outgassing from Amazonian rivers and wetlands as a large tropical source of atmospheric CO₂. *Nature* 416:617–620
- Rubio-Arias H, Quintana C, Jimenez-Castro C, Quintana R, Gutierrez M (2010) Contamination of the Conchos River in Mexico: does it pose a health risk to local residents? *Int J Environ Res Public Health* 7:2071–2084
- Salmon SD, Walter MT, Hedin LO, Brown MG (2001) Hydrological controls on chemical export from undisturbed old-growth Chilean forest. *J Hydrol* 253:69–80
- Scheren PAGM, Zanting HA, Lemmens AMC (2000) Estimation of water pollution sources in Lake Victoria, East Africa: application and elaboration of the rapid assessment methodology. *J Environ Manag* 58:235–248
- Schipper LA, Baisden WT, Parfitt RL, Ross C, Claydon JJ, Arnold G (2007) Large losses of soil C and N from soil profiles under pasture in New Zealand during the past 20 years. *Global Change Biol* 13:1138–1144
- Selva EC, Couto EG, Johnson MS, Lehmann J (2007) Litterfall production and fluvial export in headwater catchments of the southern Amazon. *J Trop Ecol* 23:329–335
- Soil Survey Staff (2003) *Keys to soil taxonomy*, 9th edn. USDA-Soil Conservation Service, Pocahontas Press, Blacksburg 332p
- Solomon D, Lehmann J, Kinyangi J, Amelung W, Lobe I, Pell A, Riha S, Ngoze S, Verchot L, Mbugua D, Skjemstad J, Schäfer T (2007) Long-term impacts of anthropogenic perturbations on dynamics and speciation of organic carbon in tropical forest and subtropical grassland ecosystems. *Global Change Biol* 13(2):511–530
- Spaans EJA, Baltissen GAM, Bouma J, Miedeme R, Lansu ALE, Schoonderbeek D, Wielemaker WG (1989) Changes in physical properties of young and old volcanic surface soils in Costa Rica after clearing of tropical rain forest. *Hydrol Process* 3:383–392
- Storer DA (1984) A simple high sample volume ashing procedure for determination of soil organic matter. *Comm Soil Sci Plant Anal* 15(7):759–772
- Stutter MI, Langan SJ, Cooper RJ (2008) Spatial and temporal dynamics of stream water particulate and dissolved N, P and C forms along a catchment transect, NE Scotland. *J Hydrol* 350:187–202
- Swallow BM, Sang JK, Nyabenge M, Bundotich DK, Duraipapp AK, Yatich TB (2009) Tradeoffs, synergies and

- traps among ecosystem services in the Lake Victoria basin of East Africa. *Environ Sci Policy* 12:504–519
- Thomas SM, Neill C, Deegan LA, Krusche AV, Ballester VM, Victoria RL (2004) Influences of land use and stream size on particulate and dissolved materials in a small Amazonian stream network. *Biogeochemistry* 68:135–151
- Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S (2002) Agricultural sustainability and intensive production practices. *Nature* 418:671–677
- Tsujimura M, Onda Y, Ito J (2001) Stream water chemistry in a steep headwater basin with high relief. *Hydrol Process* 15:1847–1858
- Veneklaas EJ (1991) Litterfall and nutrient fluxes in two montane tropical rain forests Colombia. *J Trop Ecol* 7:319–336
- Verschuren D, Johnson TC, Kling HJ, Edgington DN, Leavitt PR, Brown ET, Talbot MR, Hecky RE (2002) History and timing of human impact on Lake Victoria, East Africa. *Proc R Soc Lond B* 269:289–294
- Vidal-Abarca MR, Suarez ML, Guerrero C, Velaso J, Moreno JL, Millan A, Peran A (2001) Dynamics of dissolved and particulate organic carbon in a saline and semiarid stream of southeast Spain (Chicamo stream). *Hydrobiologia* 455:71–78
- Waldron S, Flowers H, Arlaud C, Bryant C, McFarlane S (2009) The significance of organic carbon and nutrient export from peatland-dominated landscapes subject to disturbance, a stoichiometric perspective. *Biogeosciences* 6:363–374
- Werner C, Kiese R, Butterbach-Bahl K (2007) Soil-atmosphere exchange of N₂O, CH₄, and CO₂ and controlling environmental factors for tropical rain forest sites in western Kenya. *J Geophys Res* 112(1–15):D03308
- Williams MR, Melack JM (1997) Solute export from forested and partially deforested catchments in the central Amazon. *Biogeochemistry* 38:67–102
- Williams MR, Fisher TR, Melack JM (1997) Solute dynamics in soil water and groundwater in central Amazon catchment undergoing deforestation. *Biogeochemistry* 38:303–335
- Young K, Morse GK, Scrimshaw MD, Kinniburgh JH, MacLeod CL, Lester JN (1999) The relation between phosphorus and eutrophication in the Thames catchment, UK. *Sci Total Environ* 228:157–183
- Yu J, Ho W, Lu H, Yang Y (2011) Study of water quality and genotoxicity of surface microlayer and subsurface water in Guangzhou section of Pearl River. *Environ Monit Assess* 174:681–692
- Yusop Z, Douglas I, Nik RA (2006) Export of dissolved and undissolved nutrients from forested catchments in Peninsular Malaysia. *For Ecol Manag* 224:26–44
- Zulkifli Y (1990) Effects of logging on streamwater quality and input-output budgets in small watersheds in Peninsular Malaysia. MSc. thesis, Universiti Pertanian Malaysia